

# Package: SI (via r-universe)

January 26, 2025

**Type** Package

**Title** Stochastic Integrating

**Version** 0.2.0

**Author** Jinhong Du

**Maintainer** Jinhong Du <jayduking@gmail.com>

**Description** An implementation of four stochastic methods of integrating in R, including: 1. Stochastic Point Method (or Monte Carlo Method); 2. Mean Value Method; 3. Important Sampling Method; 4. Stratified Sampling Method. It can be used to estimate one-dimension or multi-dimension integration by Monte Carlo methods. And the estimated variance (precision) is given. Reference: Caflisch, R. E. (1998) <doi:10.1017/S0962492900002804>.

**License** GPL

**Encoding** UTF-8

**LazyData** true

**Date** 2018-09-22

**Depends** R (>= 3.0.1), stats (>= 3.3.2)

**Suggests** knitr, rmarkdown, testthat

**VignetteBuilder** knitr

**RoxygenNote** 6.0.1

**NeedsCompilation** no

**Date/Publication** 2018-09-23 04:10:10 UTC

**Additional\_repositories** <https://cranhaven.r-universe.dev>

**Repository** <https://cranhaven.r-universe.dev>

**RemoteUrl** <https://github.com/cranhaven/cranhaven.r-universe.dev>

**RemoteRef** package/SI

**RemoteSha** 1fe61c0036b3e6bae85235542da8ccbc4d6f3780

**RemoteSubdir** SI

## Contents

SI.ISM . . . . .	2
SI.MVM . . . . .	3
SI.SPM . . . . .	3
SI.SSM . . . . .	4
<b>Index</b>	<b>6</b>

---

SI.ISM	<i>Important Sampling Method</i>
--------	----------------------------------

---

### Description

Important Sampling Method

### Usage

```
SI.ISM(h, g, G_inv, N, min_G = 0, max_G = 1)
```

### Arguments

h	Density function to be integrated
g	Sampling density function
G_inv	The inverse function of sampling distribution function
N	The number of trials
min_G	The min value of G
max_G	The max value of G

### Value

I	Approximated integration
Var	Estimated variance

### Examples

```
## To integrate exp(x) from -1 to 1
## Use the sampling density (3/2+x)/3
set.seed(0)
h <- function(x){
  exp(x)
}
N <- 100000
g <- function(x){return((3/2+x)/3)}
G_inv <- function(y){return(sqrt(6*y+1/4)-3/2)}
ISMresult <- SI.ISM(h,g,G_inv,N)
I3 <- ISMresult[[1]]
VarI3 <- ISMresult[[2]]
```

---

SI.MVM

*Mean Value Method*

---

**Description**

Mean Value Method

**Usage**

SI.MVM(h, from, to, N)

**Arguments**

h	Density function to be integrated
from	The start point
to	The end point
N	The number of trials

**Value**

I	Approximated integration
Var	Estimated variance

**Examples**

```
## To integrate exp(x) from -1 to 1
set.seed(0)
h <- function(x){
  exp(x)
}
N <- 100000
MVMresult <- SI.MVM(h,-1,1,N)
I2 <- MVMresult[[1]]
VarI2 <- MVMresult[[2]]
```

---

SI.SPM

*Stochastic Point Method*

---

**Description**

Stochastic Point Method

**Usage**

SI.SPM(h, from, to, M, N)

**Arguments**

h	Density function to be integrated
from	The start point
to	The end point
M	The upper bound of h(x) in [from,to]
N	The number of trials

**Value**

I	Approximated integration
Var	Estimated variance

**Examples**

```
## To integrate exp(x) from -1 to 1
set.seed(0)
h <- function(x){
  exp(x)
}
N <- 100000
SPMresult <- SI.SPM(h,-1,1,exp(1),N)
I1 <- SPMresult[[1]]
VarI1 <- SPMresult[[2]]
```

---

 SI.SSM

*Stratified Sampling Method*


---

**Description**

Stratified Sampling Method

**Usage**

```
SI.SSM(h, from, to, level, N)
```

**Arguments**

h	Density function to be integrated
from	The start point
to	The end point
level	Stratification, number of intervals
N	The number of trials

**Value**

I	Approximated integration
Var	Estimated variance

**Examples**

```
## To integrate exp(x) from -1 to 1
set.seed(0)
h <- function(x){
  exp(x)
}
N <- 100000
SSMresult <- SI.SSM(h,-1,1,10,N)
I4 <- SSMresult[[1]]
VarI4 <- SSMresult[[2]]
```

# Index

- \* **ISM**
  - SI . ISM, 2
- \* **MVM**
  - SI . MVM, 3
- \* **SPM**
  - SI . SPM, 3
  
- SI . ISM, 2
- SI . MVM, 3
- SI . SPM, 3
- SI . SSM, 4